Utah Influenza Report

This report contains data through the week ending 04/23/2016 (MMWR week 16).

UTAH DEPARTMENT OF HEALTH
Communicable Disease Analysis & Reporting Program

Overview of Influenza Surveillance: Surveillance for the 2015-2016 influenza season officially began on October 4, 2015. The Utah Department of Health publishes a weekly report throughout the active influenza season that synthesizes data from a variety of sources to give the most complete and up-to-date picture of influenza activity in the state of Utah. Data in this report should be considered provisional, and may change as more complete reports are received.

Influenza-like Illness (ILI): The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) is a national system that conducts surveillance for influenza-like illness (ILI) in outpatient healthcare facilities. ILINet providers report weekly the total number of patients seen for any reason and the number of patients seen with ILI (defined as a fever ≥ 100° F and a cough or sore throat). These data are used to determine the amount of ILI circulating in the community, as well as provide insight into regional differences in ILI activity. More than 50 facilities within 10 health jurisdictions throughout Utah participate in ILINet.

Figure 1. Weekly Influenza-like Illness Activity
Influenza-like illness (ILI) activity as measured by the number of standard deviations from the epidemic threshold - Utah, 2015-2016



Table 1. Influenza-like Illness (ILI) Activity Levels by Health District - Utah, Current Week

Health District	ILI Activity
Bear River	Minimal
Central Utah	Minimal
Davis County	Minimal
Salt Lake County	Minimal
San Juan County	No Data *
Southeast Utah	No Data *
Southwest Utah	Minimal
Summit County	Minimal
Tooele County	Minimal
TriCounty	No Data *
Utah County	Minimal
Wasatch County	Minimal
Weber-Morgan	Minimal
State Average	**Minimal**

*No participating sites in this jurisdiction

http://health.utah.gov/epi/data ● 801-538-6191 ● epi@utah.gov

Utah Influenza Report

This report contains data through the week ending 04/23/2016 (MMWR week 16).

UTAH DEPARTMENT OF HEALTH
Communicable Disease Analysis & Reporting Program

Influenza Hospitalizations: Influenza hospitalizations are a reportable condition in Utah. A person meets the case definition for an influenza hospitalization if they are hospitalized for any length of time and have an influenza positive serology, DFA, PCR, culture or rapid influenza diagnostic test. Public health in Utah gathers a variety of data on influenza hospitalizations including clinical features, course of illness, risk and protective factors, and influenza type and subtype. Data from influenza hospitalizations allows public health in Utah to better understand subgroups of the Utah population that are most severely affected by influenza and help to guide prevention messages and interventions.

Figure 2. Influenza Hospitalizations
Number of influenza hospitalizations by event date* - Utah, 2015-2016



*Event date is calculated based on a hierarchy of dates: 1. onset date 2. specimen collection date 3. date reported to public health.

Table 2. Influenza Hospitalizations by Health District - Utah

Health District	Current Week	Season To Date
Bear River	0	64
Central Utah	0	26
Davis County	2	105
Salt Lake County	8	530
San Juan County	0	0
Southeast Utah	0	3
Southwest Utah	0	96
Summit County	0	11
Tooele County	0	6
TriCounty	0	32
Utah County	2	161
Wasatch County	1	8
Weber-Morgan	3	99
State Total	**16**	**1141**

Utah Influenza Report

This report contains data through the week ending 04/23/2016 (MMWR week 16).

UTAH DEPARTMENT OF HEALTH
Communicable Disease Analysis & Reporting Program

Table 3. Influenza Hospitalizations by Age Group - Utah, Season To Date

Age Group	Total Cases	% of Cases	Rate*
0-4	120	10.5	45.2
5-24	111	9.7	11.6
25-49	203	17.8	21.3
50-64	278	24.4	67.1
65+	429	37.6	163.3
Total	**1141**	**100.0**	**40.0**

*Rate is calculated as the number of cases per 100,000 population

Table 4. Influenza Hospitalizations by Sex and Race - Utah, Season To Date

Variable		Num. of Cases	% of Cases	% in Utah Pop	p value*
Sex	Male	596	52.2	50.2	0.1792
	Female	542	47.5	49.8	0.1275
	Unknown	2	0.2	NA	--
Race	White, Not Hispanic	685	60.0	79.9	0.4425
	Hispanic	78	6.8	11.9	0.0013
	Native Hawaiian/Pacific Islander	30	2.6	1.0	<0.0001
	Black/African American	20	1.8	1.3	<0.0001
	American Indian	4	0.4	1.5	0.6433
	Asian	11	1.0	2.2	0.1644
	Unknown	313	27.4	NA	--

*If a p value is ≤ 0.05, there is a significant difference between the percentage seen in influenza hospitalizations and the general Utah population.

Utah Influenza Report

This report contains data through the week ending 04/23/2016 (MMWR week 16).

UTAH DEPARTMENT OF HEALTH
Communicable Disease Analysis & Reporting Program

Student Absenteeism: School-age children are at high risk for respiratory virus infections, including influenza. Aggregate, all-cause absenteeism data is collected weekly from over 350 schools throughout Utah. These data are analyzed to identify elevated absenteeism rates that could indicate the circulation of influenza in school-age children.

Figure 3. Student Absenteeism by Week

Rates for absences due to all causes - Utah, 2015-2016 and previous 3 year average



Table 5. Weekly Student Absenteeism - Utah, Current Week

Health District	Absences per 100 students/day
Bear River	5.0
Central Utah	5.3
Davis County	3.6
Salt Lake County	4.8
San Juan County	6.5
Southeast Utah	5.3
Southwest Utah	5.7
Summit County	3.3
Tooele County	6.1
TriCounty	6.6
Utah County	4.7
Wasatch County	3.9
Weber-Morgan	6.0
State Average	4.8

Pneumonia and Influenza Deaths: Each week the total number of death certificates received and the number of those for which pneumonia or influenza was listed as an underlying or contributing cause of death is collected. The percentage of deaths due to pneumonia and influenza are compared with a seasonal baseline and epidemic threshold value calculated for each week. These data are used to monitor the severity of influenza illness in the community.

Figure 4. Pneumonia and Influenza Deaths

Percentage of deaths attributable to pneumonia and influenza by week of death - Utah, 2010-2016



Utah Influenza Report

This report contains data through the week ending 04/23/2016 (MMWR week 16).

UTAH DEPARTMENT OF HEALTH
Communicable Disease Analysis & Reporting Program

Laboratory Surveillance: The Utah - National Electronic Disease Surveillance System (UT-NEDSS) maintains influenza testing results from hospital laboratories and the Utah Public Health Laboratory (UPHL). At UPHL, specimens are tested to determine influenza type and subtype. A portion of specimens are also sent to the Centers for Disease Control and Prevention for additional testing, including gene sequencing, antiviral resistance testing and antigenic characterization.

Figure 5. Influenza Positive Tests

Influenza positive specimens tested by laboratories throughout the state, 2015-2016



Table 6. UT-NEDSS Laboratory Influenza Testing Data: Positive Specimens by Type/Subtype

	Current Week		Season to Date	
Total Number of Positive Specimens	69		3,861	
	Number	Percentage	Number	Percentage
Influenza Type A	**38**	**55%**	**2259**	**59%**
A (2009 H1N1 Subtype)	15	39%	850	38%
A (H1 Subtype)	0	0%	33	1%
A (H3 Subtype)	6	16%	118	5%
A (No Subtyping)	17	45%	1258	56%
Influenza Type B	**31**	**45%**	**1602**	**41%**

http://health.utah.gov/epi/data ● 801-538-6191 ● epi@utah.gov