



Dear Patient:

If you are receiving this letter, your pharmacist has just dispensed to you a prescription for brand OxyContin® (oxycodone HCl controlled-release) Tablets. The purpose of this letter is to inform you that if you have had prescriptions for OxyContin filled in the past, you may have received a generic version of the medication. Therefore, your current prescription may contain tablets that look different from the medication you have previously taken.

The tablets from your current prescription for OxyContin are round in shape and the color varies depending on the strength of the tablet. OxyContin 10 mg tablets are white, 20 mg tablets are pink, 40 mg tablets are yellow, and the 80 mg tablets are green. Below is an image of OxyContin Tablets for your reference.



The sale and distribution of generic OxyContin formulations by various manufacturers is being discontinued. Some generic versions of OxyContin Tablets will continue to be available, however we cannot provide an exact timetable for when generic formulations will no longer be available.

The discontinuation of the sale of generic versions of OxyContin is not due to a safety concern. It is safe to finish your previous prescription if you still have tablets remaining.

We understand that patients are concerned about the cost of medication, and we are working to ensure that patients have appropriate access to this medication. Purdue is negotiating with managed care organizations and other insurers to encourage healthcare coverage of, and reasonable patient co-payments for, OxyContin. Furthermore, except where prohibited by state or federal law or healthcare insurers, Purdue is offering discount cards through physicians to help reduce patients' out-of-pocket costs.

If you have questions or concerns about taking OxyContin Tablets, you should talk to your doctor and pharmacist, or contact Purdue's Medical Services department at 1-888-726-7535, prompt #1. Please read the Patient Information located on back.

Sincerely yours,

A handwritten signature in cursive script that reads "Monica Kwarciński".

Monica Kwarciński, PharmD
Senior Director, Medical Services

PATIENT INFORMATION

OXYCONTIN®

(OXYCODONE HCl CONTROLLED-RELEASE) TABLETS

OxyContin® Tablets, 10 mg OxyContin® Tablets, 80 mg
OxyContin® Tablets, 20 mg OxyContin® Tablets, 160 mg
OxyContin® Tablets, 40 mg

Read this information carefully before you take OxyContin® (ox-e-CON-tin) tablets. Also read the information you get with your refills. There may be something new. This information does not take the place of talking with your doctor about your medical condition or your treatment. Only you and your doctor can decide if OxyContin is right for you. Share the important information in this leaflet with members of your household.

What Is The Most Important Information I Should Know About OxyContin®?

- Use OxyContin the way your doctor tells you to.
- Use OxyContin only for the condition for which it was prescribed.
- OxyContin is not for occasional ("as needed") use.
- Swallow the tablets whole. Do not break, crush, dissolve, or chew them before swallowing. OxyContin® works properly over 12 hours only when swallowed whole. If a tablet is broken, crushed, dissolved, or chewed, the entire 12 hour dose will be absorbed into your body all at once. This can be dangerous, causing an overdose, and possibly death.
- Keep OxyContin® out of the reach of children. Accidental overdose by a child is dangerous and may result in death.
- Prevent theft and misuse. OxyContin contains a narcotic painkiller that can be a target for people who abuse prescription medicines. Therefore, keep your tablets in a secure place, to protect them from theft. Never give them to anyone else. Selling or giving away this medicine is dangerous and against the law.

What is OxyContin®?

OxyContin® is a tablet that comes in several strengths and contains the medicine oxycodone (ox-e-KOE-done). This medicine is a painkiller like morphine. OxyContin treats moderate to severe pain that is expected to last for an extended period of time. Use OxyContin regularly during treatment. It contains enough medicine to last for up to twelve hours.

Who Should Not Take OxyContin®?

Do not take OxyContin® if

- your doctor did not prescribe OxyContin® for you.
- your pain is mild or will go away in a few days.
- your pain can be controlled by occasional use of other painkillers.
- you have severe asthma or severe lung problems.
- you have had a severe allergic reaction to codeine, hydrocodone, dihydrocodeine, or oxycodone (such as Tylox, Tylenol with Codeine, or Vicodin). A severe allergic reaction includes a severe rash, hives, breathing problems, or dizziness.
- you had surgery less than 12 - 24 hours ago and you were not taking OxyContin just before surgery.

Your doctor should know about all your medical conditions before deciding if OxyContin is right for you and what dose is best. Tell your doctor about all of your medical problems, especially the ones listed below:

- trouble breathing or lung problems
- head injury
- liver or kidney problems
- adrenal gland problems, such as Addison's disease
- convulsions or seizures
- alcoholism
- hallucinations or other severe mental problems
- past or present substance abuse or drug addiction

If any of these conditions apply to you, and you haven't told your doctor, then you should tell your doctor before taking OxyContin.

If you are pregnant or plan to become pregnant, talk with your doctor. OxyContin may not be right for you. Tell your doctor if you are breast feeding. OxyContin will pass through the milk and may harm the baby.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. They may cause serious medical problems when taken with OxyContin, especially if they cause drowsiness.

How Should I Take OxyContin®?

- Follow your doctor's directions exactly. Your doctor may change your dose based on your reactions to the medicine. Do not change your dose unless your doctor tells you to change it. Do not take OxyContin more often than prescribed.
- Swallow the tablets whole. Do not break, crush, dissolve, or chew before swallowing. If the tablets are not whole, your body will absorb too much medicine at one time. This can lead to serious problems, including overdose and death.
- If you miss a dose, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once unless your doctor tells you to.
- In case of overdose, call your local emergency number or Poison Control Center right away.
- Review your pain regularly with your doctor to determine if you still need OxyContin.
- You may see tablets in your stools (bowel movements). Do not be concerned. Your body has already absorbed the medicine.

If you continue to have pain or bothersome side effects, call your doctor.

Stopping OxyContin. Consult your doctor for instructions on how to stop this medicine slowly to avoid uncomfortable symptoms. You should not stop taking OxyContin all at once if you have been taking it for more than a few days.

After you stop taking OxyContin, flush the unused tablets down the toilet.

What Should I Avoid While Taking OxyContin®?

- Do not drive, operate heavy machinery, or participate in any other possibly dangerous activities until you know how you react to this medicine. OxyContin can make you sleepy.
- Do not drink alcohol while using OxyContin. It may increase the chance of getting dangerous side effects.
- Do not take other medicines without your doctor's approval. Other medicines include prescription and non-prescription medicines, vitamins, and supplements. Be especially careful about products that make you sleepy.

What are the Possible Side Effects of OxyContin®?

Call your doctor or get medical help right away if

- your breathing slows down
 - you feel faint, dizzy, confused, or have any other unusual symptoms
- Some of the common side effects of OxyContin® are nausea, vomiting, dizziness, drowsiness, constipation, itching, dry mouth, sweating, weakness, and headache. Some of these side effects may decrease with continued use.

There is a risk of abuse or addiction with narcotic painkillers. If you have abused drugs in the past, you may have a higher chance of developing abuse or addiction again while using OxyContin.

These are not all the possible side effects of OxyContin. For a complete list, ask your doctor or pharmacist.

General Advice About OxyContin

- Do not use OxyContin for conditions for which it was not prescribed.
- Do not give OxyContin to other people, even if they have the same symptoms you have. Sharing is illegal and may cause severe medical problems, including death.

This leaflet summarizes the most important information about OxyContin. If you would like more information, talk with your doctor. Also, you can ask your pharmacist or doctor for information about OxyContin that is written for health professionals.

Rx Only

©2002, 2004, 2005, 2007, Purdue Pharma L.P.
Purdue Pharma L.P., Stamford, CT 06901-3431
January 15, 2007 0T00367J